



AP[®] Calculus BC 2004 Scoring Guidelines

The materials included in these files are intended for noncommercial use by AP teachers for course and exam preparation; permission for any other use must be sought from the Advanced Placement Program[®]. Teachers may reproduce them, in whole or in part, in limited quantities, for face-to-face teaching purposes but may not mass distribute the materials, electronically or otherwise. This permission does not apply to any third-party copyrights contained herein. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here.

The College Board is a not-for-profit membership association whose mission is to connect students to college success and opportunity. Founded in 1900, the association is composed of more than 4,500 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 23,000 high schools, and 3,500 colleges through major programs and services in college admissions, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT[®], the PSAT/NMSQT[®], and the Advanced Placement Program[®] (AP[®]). The College Board is committed to the principles of excellence and equity, and that commitment is embodied in all of its programs, services, activities, and concerns.

For further information, visit www.collegeboard.com

Copyright © 2004 College Entrance Examination Board. All rights reserved. College Board, Advanced Placement Program, AP, AP Central, AP Vertical Teams, APCD, Pacesetter, Pre-AP, SAT, Student Search Service, and the acorn logo are registered trademarks of the College Entrance Examination Board. PSAT/NMSQT is a registered trademark of the College Entrance Examination Board and National Merit Scholarship Corporation. Educational Testing Service and ETS are registered trademarks of Educational Testing Service. Other products and services may be trademarks of their respective owners.

For the College Board's online home for AP professionals, visit AP Central at apcentral.collegeboard.com.

AP[®] CALCULUS BC
2004 SCORING GUIDELINES

Question 1

Traffic flow is defined as the rate at which cars pass through an intersection, measured in cars per minute. The traffic flow at a particular intersection is modeled by the function F defined by

$$F(t) = 82 + 4\sin\left(\frac{t}{2}\right) \text{ for } 0 \leq t \leq 30,$$

where $F(t)$ is measured in cars per minute and t is measured in minutes.

- (a) To the nearest whole number, how many cars pass through the intersection over the 30-minute period?
- (b) Is the traffic flow increasing or decreasing at $t = 7$? Give a reason for your answer.
- (c) What is the average value of the traffic flow over the time interval $10 \leq t \leq 15$? Indicate units of measure.
- (d) What is the average rate of change of the traffic flow over the time interval $10 \leq t \leq 15$? Indicate units of measure.

(a) $\int_0^{30} F(t) dt = 2474$ cars

3 : $\begin{cases} 1 : \text{limits} \\ 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(b) $F'(7) = -1.872$ or -1.873
Since $F'(7) < 0$, the traffic flow is decreasing at $t = 7$.

1 : answer with reason

(c) $\frac{1}{5} \int_{10}^{15} F(t) dt = 81.899$ cars/min

3 : $\begin{cases} 1 : \text{limits} \\ 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(d) $\frac{F(15) - F(10)}{15 - 10} = 1.517$ or 1.518 cars/min²

1 : answer


Units of cars/min in (c) and cars/min² in (d)

1 : units in (c) and (d)

**AP[®] CALCULUS BC
2004 SCORING GUIDELINES**

Question 2

Let f and g be the functions given by $f(x) = 2x(1 - x)$ and $g(x) = 3(x - 1)\sqrt{x}$ for $0 \leq x \leq 1$. The graphs of f and g are shown in the figure above.



- (a) Find the area of the shaded region enclosed by the graphs of f and g .
- (b) Find the volume of the solid generated when the shaded region enclosed by the graphs of f and g is revolved about the horizontal line $y = 2$.
- (c) Let h be the function given by $h(x) = kx(1 - x)$ for $0 \leq x \leq 1$. For each $k > 0$, the region (not shown) enclosed by the graphs of h and g is the base of a solid with square cross sections perpendicular to the x -axis. There is a value of k for which the volume of this solid is equal to 15. Write, but do not solve, an equation involving an integral expression that could be used to find the value of k .

(a) Area = $\int_0^1 (f(x) - g(x)) dx$
 $= \int_0^1 (2x(1 - x) - 3(x - 1)\sqrt{x}) dx = 1.133$

2 : $\left\{ \begin{array}{l} 1 : \text{integral} \\ 1 : \text{answer} \end{array} \right.$

(b) Volume = $\pi \int_0^1 ((2 - g(x))^2 - (2 - f(x))^2) dx$
 $= \pi \int_0^1 ((2 - 3(x - 1)\sqrt{x})^2 - (2 - 2x(1 - x))^2) dx$
 $= 16.179$

4 : $\left\{ \begin{array}{l} 1 : \text{limits and constant} \\ 2 : \text{integrand} \\ \langle -1 \rangle \text{ each error} \\ \text{Note: } 0/2 \text{ if integral not of form} \\ \int_a^b (R^2(x) - r^2(x)) dx \\ 1 : \text{answer} \end{array} \right.$

(c) Volume = $\int_0^1 (h(x) - g(x))^2 dx$
 $\int_0^1 (kx(1 - x) - 3(x - 1)\sqrt{x})^2 dx = 15$

3 : $\left\{ \begin{array}{l} 2 : \text{integrand} \\ 1 : \text{answer} \end{array} \right.$

**AP[®] CALCULUS BC
2004 SCORING GUIDELINES**

Question 3

An object moving along a curve in the xy -plane has position $(x(t), y(t))$ at time $t \geq 0$ with $\frac{dx}{dt} = 3 + \cos(t^2)$. The derivative $\frac{dy}{dt}$ is not explicitly given. At time $t = 2$, the object is at position $(1, 8)$.

- (a) Find the x -coordinate of the position of the object at time $t = 4$.
- (b) At time $t = 2$, the value of $\frac{dy}{dt}$ is -7 . Write an equation for the line tangent to the curve at the point $(x(2), y(2))$.
- (c) Find the speed of the object at time $t = 2$.
- (d) For $t \geq 3$, the line tangent to the curve at $(x(t), y(t))$ has a slope of $2t + 1$. Find the acceleration vector of the object at time $t = 4$.

(a)
$$x(4) = x(2) + \int_2^4 (3 + \cos(t^2)) dt$$

$$= 1 + \int_2^4 (3 + \cos(t^2)) dt = 7.132 \text{ or } 7.133$$

3 :
$$\begin{cases} 1 : \int_2^4 (3 + \cos(t^2)) dt \\ 1 : \text{handles initial condition} \\ 1 : \text{answer} \end{cases}$$

(b)
$$\left. \frac{dy}{dx} \right|_{t=2} = \left. \frac{\frac{dy}{dt}}{\frac{dx}{dt}} \right|_{t=2} = \frac{-7}{3 + \cos 4} = -2.983$$

$$y - 8 = -2.983(x - 1)$$

2 :
$$\begin{cases} 1 : \text{finds } \left. \frac{dy}{dx} \right|_{t=2} \\ 1 : \text{equation} \end{cases}$$

(c) The speed of the object at time $t = 2$ is $\sqrt{(x'(2))^2 + (y'(2))^2} = 7.382 \text{ or } 7.383$.

1 : answer

(d) $x''(4) = 2.303$

$$y'(t) = \frac{dy}{dt} = \frac{dy}{dx} \cdot \frac{dx}{dt} = (2t + 1)(3 + \cos(t^2))$$

$$y''(4) = 24.813 \text{ or } 24.814$$

The acceleration vector at $t = 4$ is $\langle 2.303, 24.813 \rangle$ or $\langle 2.303, 24.814 \rangle$.

3 :
$$\begin{cases} 1 : x''(4) \\ 1 : \frac{dy}{dt} \\ 1 : \text{answer} \end{cases}$$

**AP[®] CALCULUS BC
2004 SCORING GUIDELINES**

Question 4

Consider the curve given by $x^2 + 4y^2 = 7 + 3xy$.

- (a) Show that $\frac{dy}{dx} = \frac{3y - 2x}{8y - 3x}$.
- (b) Show that there is a point P with x -coordinate 3 at which the line tangent to the curve at P is horizontal. Find the y -coordinate of P .
- (c) Find the value of $\frac{d^2y}{dx^2}$ at the point P found in part (b). Does the curve have a local maximum, a local minimum, or neither at the point P ? Justify your answer.

(a)
$$\begin{aligned} 2x + 8yy' &= 3y + 3xy' \\ (8y - 3x)y' &= 3y - 2x \\ y' &= \frac{3y - 2x}{8y - 3x} \end{aligned}$$

2 : $\begin{cases} 1 : \text{implicit differentiation} \\ 1 : \text{solves for } y' \end{cases}$

(b) $\frac{3y - 2x}{8y - 3x} = 0; 3y - 2x = 0$

When $x = 3$, $3y = 6$
 $y = 2$

$3^2 + 4 \cdot 2^2 = 25$ and $7 + 3 \cdot 3 \cdot 2 = 25$

Therefore, $P = (3, 2)$ is on the curve and the slope is 0 at this point.

3 : $\begin{cases} 1 : \frac{dy}{dx} = 0 \\ 1 : \text{shows slope is 0 at } (3, 2) \\ 1 : \text{shows } (3, 2) \text{ lies on curve} \end{cases}$

(c)
$$\frac{d^2y}{dx^2} = \frac{(8y - 3x)(3y' - 2) - (3y - 2x)(8y' - 3)}{(8y - 3x)^2}$$

At $P = (3, 2)$, $\frac{d^2y}{dx^2} = \frac{(16 - 9)(-2) - (-2)(8 - 3)}{(16 - 9)^2} = -\frac{2}{7}$.

Since $y' = 0$ and $y'' < 0$ at P , the curve has a local maximum at P .

4 : $\begin{cases} 2 : \frac{d^2y}{dx^2} \\ 1 : \text{value of } \frac{d^2y}{dx^2} \text{ at } (3, 2) \\ 1 : \text{conclusion with justification} \end{cases}$

**AP[®] CALCULUS BC
2004 SCORING GUIDELINES**

Question 5

A population is modeled by a function P that satisfies the logistic differential equation

$$\frac{dP}{dt} = \frac{P}{5} \left(1 - \frac{P}{12} \right).$$

(a) If $P(0) = 3$, what is $\lim_{t \rightarrow \infty} P(t)$?

If $P(0) = 20$, what is $\lim_{t \rightarrow \infty} P(t)$?

(b) If $P(0) = 3$, for what value of P is the population growing the fastest?

(c) A different population is modeled by a function Y that satisfies the separable differential equation

$$\frac{dY}{dt} = \frac{Y}{5} \left(1 - \frac{t}{12} \right).$$

Find $Y(t)$ if $Y(0) = 3$.

(d) For the function Y found in part (c), what is $\lim_{t \rightarrow \infty} Y(t)$?

(a) For this logistic differential equation, the carrying capacity is 12.

If $P(0) = 3$, $\lim_{t \rightarrow \infty} P(t) = 12$.

If $P(0) = 20$, $\lim_{t \rightarrow \infty} P(t) = 12$.

2 : $\begin{cases} 1 : \text{answer} \\ 1 : \text{answer} \end{cases}$

(b) The population is growing the fastest when P is half the carrying capacity. Therefore, P is growing the fastest when $P = 6$.

1 : answer

(c) $\frac{1}{Y} dY = \frac{1}{5} \left(1 - \frac{t}{12} \right) dt = \left(\frac{1}{5} - \frac{t}{60} \right) dt$

$$\ln|Y| = \frac{t}{5} - \frac{t^2}{120} + C$$

$$Y(t) = K e^{\frac{t}{5} - \frac{t^2}{120}}$$

$$K = 3$$

$$Y(t) = 3e^{\frac{t}{5} - \frac{t^2}{120}}$$

5 : $\begin{cases} 1 : \text{separates variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } Y \\ 0/1 \text{ if } Y \text{ is not exponential} \end{cases}$

Note: max 2/5 [1-1-0-0-0] if no constant of integration

Note: 0/5 if no separation of variables

(d) $\lim_{t \rightarrow \infty} Y(t) = 0$

1 : answer
0/1 if Y is not exponential

AP[®] CALCULUS BC
2004 SCORING GUIDELINES

Question 6

Let f be the function given by $f(x) = \sin\left(5x + \frac{\pi}{4}\right)$, and let $P(x)$ be the third-degree Taylor polynomial for f about $x = 0$.

- (a) Find $P(x)$.
- (b) Find the coefficient of x^{22} in the Taylor series for f about $x = 0$.
- (c) Use the Lagrange error bound to show that $\left|f\left(\frac{1}{10}\right) - P\left(\frac{1}{10}\right)\right| < \frac{1}{100}$.
- (d) Let G be the function given by $G(x) = \int_0^x f(t) dt$. Write the third-degree Taylor polynomial for G about $x = 0$.

<p>(a) $f(0) = \sin\left(\frac{\pi}{4}\right) = \frac{\sqrt{2}}{2}$ $f'(0) = 5\cos\left(\frac{\pi}{4}\right) = \frac{5\sqrt{2}}{2}$ $f''(0) = -25\sin\left(\frac{\pi}{4}\right) = -\frac{25\sqrt{2}}{2}$ $f'''(0) = -125\cos\left(\frac{\pi}{4}\right) = -\frac{125\sqrt{2}}{2}$ $P(x) = \frac{\sqrt{2}}{2} + \frac{5\sqrt{2}}{2}x - \frac{25\sqrt{2}}{2(2!)}x^2 - \frac{125\sqrt{2}}{2(3!)}x^3$</p>	<p>4 : $P(x)$</p> <p>$\langle -1 \rangle$ each error or missing term</p> <p>deduct only once for $\sin\left(\frac{\pi}{4}\right)$ evaluation error</p> <p>deduct only once for $\cos\left(\frac{\pi}{4}\right)$ evaluation error</p> <p>$\langle -1 \rangle$ max for all extra terms, + \dots, misuse of equality</p>
<p>(b) $\frac{-5^{22}\sqrt{2}}{2(22!)}$</p>	<p>2 : $\begin{cases} 1 : \text{magnitude} \\ 1 : \text{sign} \end{cases}$</p>
<p>(c) $\left f\left(\frac{1}{10}\right) - P\left(\frac{1}{10}\right)\right \leq \max_{0 \leq c \leq \frac{1}{10}} f^{(4)}(c) \left(\frac{1}{4!}\right) \left(\frac{1}{10}\right)^4$ $\leq \frac{625}{4!} \left(\frac{1}{10}\right)^4 = \frac{1}{384} < \frac{1}{100}$</p>	<p>1 : error bound in an appropriate inequality</p>
<p>(d) The third-degree Taylor polynomial for G about $x = 0$ is $\int_0^x \left(\frac{\sqrt{2}}{2} + \frac{5\sqrt{2}}{2}t - \frac{25\sqrt{2}}{4}t^2\right) dt$ $= \frac{\sqrt{2}}{2}x + \frac{5\sqrt{2}}{4}x^2 - \frac{25\sqrt{2}}{12}x^3$</p>	<p>2 : third-degree Taylor polynomial for G about $x = 0$</p> <p>$\langle -1 \rangle$ each incorrect or missing term</p> <p>$\langle -1 \rangle$ max for all extra terms, + \dots, misuse of equality</p>