



## AP<sup>®</sup> Calculus BC 2006 Scoring Guidelines

### **The College Board: Connecting Students to College Success**

The College Board is a not-for-profit membership association whose mission is to connect students to college success and opportunity. Founded in 1900, the association is composed of more than 5,000 schools, colleges, universities, and other educational organizations. Each year, the College Board serves seven million students and their parents, 23,000 high schools, and 3,500 colleges through major programs and services in college admissions, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT<sup>®</sup>, the PSAT/NMSQT<sup>®</sup>, and the Advanced Placement Program<sup>®</sup> (AP<sup>®</sup>). The College Board is committed to the principles of excellence and equity, and that commitment is embodied in all of its programs, services, activities, and concerns.

© 2006 The College Board. All rights reserved. College Board, AP Central, APCD, Advanced Placement Program, AP, AP Vertical Teams, Pre-AP, SAT, and the acorn logo are registered trademarks of the College Board. Admitted Class Evaluation Service, CollegeEd, connect to college success, MyRoad, SAT Professional Development, SAT Readiness Program, and Setting the Cornerstones are trademarks owned by the College Board. PSAT/NMSQT is a registered trademark of the College Board and National Merit Scholarship Corporation. All other products and services may be trademarks of their respective owners. Permission to use copyrighted College Board materials may be requested online at: [www.collegeboard.com/inquiry/cbpermit.html](http://www.collegeboard.com/inquiry/cbpermit.html).

**Visit the College Board on the Web: [www.collegeboard.com](http://www.collegeboard.com).**


**AP Central is the official online home for the AP Program: [apcentral.collegeboard.com](http://apcentral.collegeboard.com).**

**AP<sup>®</sup> CALCULUS BC  
2006 SCORING GUIDELINES**

**Question 1**

Let  $R$  be the shaded region bounded by the graph of  $y = \ln x$  and the line  $y = x - 2$ , as shown above.

- (a) Find the area of  $R$ .
- (b) Find the volume of the solid generated when  $R$  is rotated about the horizontal line  $y = -3$ .
- (c) Write, but do not evaluate, an integral expression that can be used to find the volume of the solid generated when  $R$  is rotated about the  $y$ -axis.



$\ln(x) = x - 2$  when  $x = 0.15859$  and  $3.14619$ .  
Let  $S = 0.15859$  and  $T = 3.14619$

(a) Area of  $R = \int_S^T (\ln(x) - (x - 2)) dx = 1.949$

3 : { 1 : integrand  
1 : limits  
1 : answer

(b) Volume =  $\pi \int_S^T ((\ln(x) + 3)^2 - (x - 2 + 3)^2) dx$   
= 34.198 or 34.199

3 : { 2 : integrand  
1 : limits, constant, and answer


(c) Volume =  $\pi \int_{S-2}^{T-2} ((y + 2)^2 - (e^y)^2) dy$

3 : { 2 : integrand  
1 : limits and constant

**AP<sup>®</sup> CALCULUS BC  
2006 SCORING GUIDELINES**

**Question 2**

At an intersection in Thomasville, Oregon, cars turn left at the rate  $L(t) = 60\sqrt{t} \sin^2\left(\frac{t}{3}\right)$  cars per hour over the time interval  $0 \leq t \leq 18$  hours. The graph of  $y = L(t)$  is shown above.



- (a) To the nearest whole number, find the total number of cars turning left at the intersection over the time interval  $0 \leq t \leq 18$  hours.
- (b) Traffic engineers will consider turn restrictions when  $L(t) \geq 150$  cars per hour. Find all values of  $t$  for which  $L(t) \geq 150$  and compute the average value of  $L$  over this time interval. Indicate units of measure.
- (c) Traffic engineers will install a signal if there is any two-hour time interval during which the product of the total number of cars turning left and the total number of oncoming cars traveling straight through the intersection is greater than 200,000. In every two-hour time interval, 500 oncoming cars travel straight through the intersection. Does this intersection require a traffic signal? Explain the reasoning that leads to your conclusion.

(a)  $\int_0^{18} L(t) dt \approx 1658$  cars

(b)  $L(t) = 150$  when  $t = 12.42831, 16.12166$   
 Let  $R = 12.42831$  and  $S = 16.12166$   
 $L(t) \geq 150$  for  $t$  in the interval  $[R, S]$

$$\frac{1}{S - R} \int_R^S L(t) dt = 199.426 \text{ cars per hour}$$

- (c) For the product to exceed 200,000, the number of cars turning left in a two-hour interval must be greater than 400.

$$\int_{13}^{15} L(t) dt = 431.931 > 400$$

OR

The number of cars turning left will be greater than 400 on a two-hour interval if  $L(t) \geq 200$  on that interval.

$$L(t) \geq 200 \text{ on any two-hour subinterval of } [13.25304, 15.32386].$$

Yes, a traffic signal is required.

$$2 : \begin{cases} 1 : \text{setup} \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 1 : t\text{-interval when } L(t) \geq 150 \\ 1 : \text{average value integral} \\ 1 : \text{answer with units} \end{cases}$$

$$4 : \begin{cases} 1 : \text{considers 400 cars} \\ 1 : \text{valid interval } [h, h + 2] \\ 1 : \text{value of } \int_h^{h+2} L(t) dt \\ 1 : \text{answer and explanation} \end{cases}$$

OR

$$4 : \begin{cases} 1 : \text{considers 200 cars per hour} \\ 1 : \text{solves } L(t) \geq 200 \\ 1 : \text{discusses 2 hour interval} \\ 1 : \text{answer and explanation} \end{cases}$$

**AP<sup>®</sup> CALCULUS BC  
2006 SCORING GUIDELINES**

**Question 3**

An object moving along a curve in the  $xy$ -plane is at position  $(x(t), y(t))$  at time  $t$ , where

$$\frac{dx}{dt} = \sin^{-1}(1 - 2e^{-t}) \quad \text{and} \quad \frac{dy}{dt} = \frac{4t}{1 + t^3}$$

for  $t \geq 0$ . At time  $t = 2$ , the object is at the point  $(6, -3)$ . (Note:  $\sin^{-1}x = \arcsin x$ )

- (a) Find the acceleration vector and the speed of the object at time  $t = 2$ .  
 (b) The curve has a vertical tangent line at one point. At what time  $t$  is the object at this point?  
 (c) Let  $m(t)$  denote the slope of the line tangent to the curve at the point  $(x(t), y(t))$ . Write an expression for  $m(t)$  in terms of  $t$  and use it to evaluate  $\lim_{t \rightarrow \infty} m(t)$ .  
 (d) The graph of the curve has a horizontal asymptote  $y = c$ . Write, but do not evaluate, an expression involving an improper integral that represents this value  $c$ .

(a)  $a(2) = \langle 0.395 \text{ or } 0.396, -0.741 \text{ or } -0.740 \rangle$   
 Speed  $\sqrt{x'(2)^2 + y'(2)^2} \quad 1.207 \text{ or } 1.208$

2 :  $\begin{cases} 1 : \text{acceleration} \\ 1 : \text{speed} \end{cases}$

(b)  $\sin^{-1}(1 - 2e^{-t}) = 0$   
 $1 - 2e^{-t} = 0$   
 $t = \ln 2 = 0.693$  and  $\frac{dy}{dt} \neq 0$  when  $t = \ln 2$

2 :  $\begin{cases} 1 : x'(t) = 0 \\ 1 : \text{answer} \end{cases}$

(c)  $m(t) = \frac{4t}{1 + t^3} \cdot \frac{1}{\sin^{-1}(1 - 2e^{-t})}$   
 $\lim_{t \rightarrow \infty} m(t) = \lim_{t \rightarrow \infty} \left( \frac{4t}{1 + t^3} \cdot \frac{1}{\sin^{-1}(1 - 2e^{-t})} \right)$   
 $= 0 \left( \frac{1}{\sin^{-1}(1)} \right) = 0$

2 :  $\begin{cases} 1 : m(t) \\ 1 : \text{limit value} \end{cases}$

(d) Since  $\lim_{t \rightarrow \infty} x(t) = \infty$ ,  
 $c = \lim_{t \rightarrow \infty} y(t) = -3 + \int_2^{\infty} \frac{4t}{1 + t^3} dt$

3 :  $\begin{cases} 1 : \text{integrand} \\ 1 : \text{limits} \\ 1 : \text{initial value consistent} \\ \quad \text{with lower limit} \end{cases}$

**AP<sup>®</sup> CALCULUS BC  
2006 SCORING GUIDELINES**

**Question 4**

$t$ (seconds)	0	10	20	30	40	50	60	70	80
$v(t)$ (feet per second)	5	14	22	29	35	40	44	47	49

Rocket  $A$  has positive velocity  $v(t)$  after being launched upward from an initial height of 0 feet at time  $t = 0$  seconds. The velocity of the rocket is recorded for selected values of  $t$  over the interval  $0 \leq t \leq 80$  seconds, as shown in the table above.

(a) Find the average acceleration of rocket  $A$  over the time interval  $0 \leq t \leq 80$  seconds. Indicate units of measure.

(b) Using correct units, explain the meaning of  $\int_{10}^{70} v(t) dt$  in terms of the rocket's flight. Use a midpoint Riemann sum with 3 subintervals of equal length to approximate  $\int_{10}^{70} v(t) dt$ .

(c) Rocket  $B$  is launched upward with an acceleration of  $a(t) = \frac{3}{\sqrt{t+1}}$  feet per second per second. At time  $t = 0$  seconds, the initial height of the rocket is 0 feet, and the initial velocity is 2 feet per second. Which of the two rockets is traveling faster at time  $t = 80$  seconds? Explain your answer.

(a) Average acceleration of rocket  $A$  is

$$\frac{v(80) - v(0)}{80 - 0} = \frac{49 - 5}{80} = \frac{11}{20} \text{ ft/sec}^2$$

(b) Since the velocity is positive,  $\int_{10}^{70} v(t) dt$  represents the distance, in feet, traveled by rocket  $A$  from  $t = 10$  seconds to  $t = 70$  seconds.

A midpoint Riemann sum is

$$20[v(20) + v(40) + v(60)]$$

$$= 20[22 + 35 + 44] = 2020 \text{ ft}$$

(c) Let  $v_B(t)$  be the velocity of rocket  $B$  at time  $t$ .

$$v_B(t) = \int \frac{3}{\sqrt{t+1}} dt = 6\sqrt{t+1} + C$$

$$2 = v_B(0) = 6 + C$$

$$v_B(t) = 6\sqrt{t+1} - 4$$

$$v_B(80) = 50 > 49 = v(80)$$

Rocket  $B$  is traveling faster at time  $t = 80$  seconds.

Units of  $\text{ft/sec}^2$  in (a) and ft in (b)

1 : answer

3 :  $\left\{ \begin{array}{l} 1 : \text{explanation} \\ 1 : \text{uses } v(20), v(40), v(60) \\ 1 : \text{value} \end{array} \right.$

4 :  $\left\{ \begin{array}{l} 1 : 6\sqrt{t+1} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{finds } v_B(80), \text{ compares to } v(80), \\ \text{and draws a conclusion} \end{array} \right.$

1 : units in (a) and (b)

**AP<sup>®</sup> CALCULUS BC  
2006 SCORING GUIDELINES**

**Question 5**

Consider the differential equation  $\frac{dy}{dx} = 5x^2 - \frac{6}{y-2}$  for  $y \neq 2$ . Let  $y = f(x)$  be the particular solution to this differential equation with the initial condition  $f(-1) = -4$ .

- (a) Evaluate  $\frac{dy}{dx}$  and  $\frac{d^2y}{dx^2}$  at  $(-1, -4)$ .
- (b) Is it possible for the  $x$ -axis to be tangent to the graph of  $f$  at some point? Explain why or why not.
- (c) Find the second-degree Taylor polynomial for  $f$  about  $x = -1$ .
- (d) Use Euler's method, starting at  $x = -1$  with two steps of equal size, to approximate  $f(0)$ . Show the work that leads to your answer.

(a)  $\frac{dy}{dx} \Big|_{(-1, -4)} = 6$

$$\frac{d^2y}{dx^2} = 10x + 6(y-2)^{-2} \frac{dy}{dx}$$

$$\frac{d^2y}{dx^2} \Big|_{(-1, -4)} = -10 + 6 \frac{1}{(-6)^2} 6 = -9$$

(b) The  $x$ -axis will be tangent to the graph of  $f$  if  $\frac{dy}{dx} \Big|_{(k, 0)} = 0$ .

The  $x$ -axis will never be tangent to the graph of  $f$  because

$$\frac{dy}{dx} \Big|_{(k, 0)} = 5k^2 + 3 > 0 \text{ for all } k.$$

(c)  $P(x) = -4 + 6(x+1) - \frac{9}{2}(x+1)^2$

(d)  $f(-1) = -4$

$$f\left(-\frac{1}{2}\right) \approx -4 + \frac{1}{2}(6) = -1$$

$$f(0) \approx -1 + \frac{1}{2}\left(\frac{5}{4} + 2\right) = \frac{5}{8}$$

$$3 : \begin{cases} 1 : \frac{dy}{dx} \Big|_{(-1, -4)} \\ 1 : \frac{d^2y}{dx^2} \\ 1 : \frac{d^2y}{dx^2} \Big|_{(-1, -4)} \end{cases}$$

$$2 : \begin{cases} 1 : \frac{dy}{dx} = 0 \text{ and } y = 0 \\ 1 : \text{answer and explanation} \end{cases}$$

$$2 : \begin{cases} 1 : \text{quadratic and centered at } x = -1 \\ 1 : \text{coefficients} \end{cases}$$

$$2 : \begin{cases} 1 : \text{Euler's method with 2 steps} \\ 1 : \text{Euler's approximation to } f(0) \end{cases}$$

**AP<sup>®</sup> CALCULUS BC  
2006 SCORING GUIDELINES**

**Question 6**

The function  $f$  is defined by the power series

$$f(x) = -\frac{x}{2} + \frac{2x^2}{3} - \frac{3x^3}{4} + \cdots + \frac{(-1)^n nx^n}{n+1} + \cdots$$

for all real numbers  $x$  for which the series converges. The function  $g$  is defined by the power series

$$g(x) = 1 - \frac{x}{2!} + \frac{x^2}{4!} - \frac{x^3}{6!} + \cdots + \frac{(-1)^n x^n}{(2n)!} + \cdots$$

for all real numbers  $x$  for which the series converges.

- (a) Find the interval of convergence of the power series for  $f$ . Justify your answer.
- (b) The graph of  $y = f(x) - g(x)$  passes through the point  $(0, -1)$ . Find  $y'(0)$  and  $y''(0)$ . Determine whether  $y$  has a relative minimum, a relative maximum, or neither at  $x = 0$ . Give a reason for your answer.

(a) 
$$\left| \frac{(-1)^{n+1} (n+1)x^{n+1}}{n+2} \cdot \frac{n+1}{(-1)^n nx^n} \right| = \frac{(n+1)^2}{(n+2)(n)} \cdot |x|$$

$$\lim_{n \rightarrow \infty} \frac{(n+1)^2}{(n+2)(n)} \cdot |x| = |x|$$

The series converges when  $-1 < x < 1$ .

When  $x = 1$ , the series is  $-\frac{1}{2} + \frac{2}{3} - \frac{3}{4} + \cdots$

This series does not converge, because the limit of the individual terms is not zero.

When  $x = -1$ , the series is  $\frac{1}{2} + \frac{2}{3} + \frac{3}{4} + \cdots$

This series does not converge, because the limit of the individual terms is not zero.

Thus, the interval of convergence is  $-1 < x < 1$ .

(b)  $f'(x) = -\frac{1}{2} + \frac{4}{3}x - \frac{9}{4}x^2 + \cdots$  and  $f'(0) = -\frac{1}{2}$ .

$$g'(x) = -\frac{1}{2!} + \frac{2}{4!}x - \frac{3}{6!}x^2 + \cdots$$
 and  $g'(0) = -\frac{1}{2}$ .

$$y'(0) = f'(0) - g'(0) = 0$$

$$f''(0) = \frac{4}{3} \text{ and } g''(0) = \frac{2}{4!} = \frac{1}{12}$$

$$\text{Thus, } y''(0) = \frac{4}{3} - \frac{1}{12} > 0.$$

Since  $y'(0) = 0$  and  $y''(0) > 0$ ,  $y$  has a relative minimum at  $x = 0$ .

5 : {  
 1 : sets up ratio  
 1 : computes limit of ratio  
 1 : identifies radius of convergence  
 1 : considers both endpoints  
 1 : analysis/conclusion for both endpoints

4 : {  
 1 :  $y'(0)$   
 1 :  $y''(0)$   
 1 : conclusion  
 1 : reasoning