



**AP<sup>®</sup> Calculus BC**  
**2007 Free-Response Questions**  
**Form B**

**The College Board: Connecting Students to College Success**

The College Board is a not-for-profit membership association whose mission is to connect students to college success and opportunity. Founded in 1900, the association is composed of more than 5,000 schools, colleges, universities, and other educational organizations. Each year, the College Board serves seven million students and their parents, 23,000 high schools, and 3,500 colleges through major programs and services in college admissions, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT<sup>®</sup>, the PSAT/NMSQT<sup>®</sup>, and the Advanced Placement Program<sup>®</sup> (AP<sup>®</sup>). The College Board is committed to the principles of excellence and equity, and that commitment is embodied in all of its programs, services, activities, and concerns.

© 2007 The College Board. All rights reserved. College Board, Advanced Placement Program, AP, AP Central, SAT, and the acorn logo are registered trademarks of the College Board. PSAT/NMSQT is a registered trademark of the College Board and National Merit Scholarship Corporation.

Permission to use copyrighted College Board materials may be requested online at:  
[www.collegeboard.com/inquiry/cbpermit.html](http://www.collegeboard.com/inquiry/cbpermit.html).


Visit the College Board on the Web: [www.collegeboard.com](http://www.collegeboard.com).

AP Central is the official online home for the AP Program: [apcentral.collegeboard.com](http://apcentral.collegeboard.com).

CALCULUS BC  
SECTION II, Part A  
Time—45 minutes  
Number of problems—3

A graphing calculator is required for some problems or parts of problems.

---



1. Let  $R$  be the region bounded by the graph of  $y = e^{2x-x^2}$  and the horizontal line  $y = 2$ , and let  $S$  be the region bounded by the graph of  $y = e^{2x-x^2}$  and the horizontal lines  $y = 1$  and  $y = 2$ , as shown above.
- (a) Find the area of  $R$ .
- (b) Find the area of  $S$ .
- (c) Write, but do not evaluate, an integral expression that gives the volume of the solid generated when  $R$  is rotated about the horizontal line  $y = 1$ .
- 

**WRITE ALL WORK IN THE EXAM BOOKLET.**

2. An object moving along a curve in the  $xy$ -plane is at position  $(x(t), y(t))$  at time  $t$  with

$$\frac{dx}{dt} = \arctan\left(\frac{t}{1+t}\right) \text{ and } \frac{dy}{dt} = \ln(t^2 + 1)$$

for  $t \geq 0$ . At time  $t = 0$ , the object is at position  $(-3, -4)$ . (Note:  $\tan^{-1} x = \arctan x$ )

- Find the speed of the object at time  $t = 4$ .
  - Find the total distance traveled by the object over the time interval  $0 \leq t \leq 4$ .
  - Find  $x(4)$ .
  - For  $t > 0$ , there is a point on the curve where the line tangent to the curve has slope 2. At what time  $t$  is the object at this point? Find the acceleration vector at this point.
- 

**WRITE ALL WORK IN THE EXAM BOOKLET.**


3. The wind chill is the temperature, in degrees Fahrenheit ( $^{\circ}\text{F}$ ), a human feels based on the air temperature, in degrees Fahrenheit, and the wind velocity  $v$ , in miles per hour (mph). If the air temperature is  $32^{\circ}\text{F}$ , then the wind chill is given by  $W(v) = 55.6 - 22.1v^{0.16}$  and is valid for  $5 \leq v \leq 60$ .
- (a) Find  $W'(20)$ . Using correct units, explain the meaning of  $W'(20)$  in terms of the wind chill.
- (b) Find the average rate of change of  $W$  over the interval  $5 \leq v \leq 60$ . Find the value of  $v$  at which the instantaneous rate of change of  $W$  is equal to the average rate of change of  $W$  over the interval  $5 \leq v \leq 60$ .
- (c) Over the time interval  $0 \leq t \leq 4$  hours, the air temperature is a constant  $32^{\circ}\text{F}$ . At time  $t = 0$ , the wind velocity is  $v = 20$  mph. If the wind velocity increases at a constant rate of 5 mph per hour, what is the rate of change of the wind chill with respect to time at  $t = 3$  hours? Indicate units of measure.
- 

**WRITE ALL WORK IN THE EXAM BOOKLET.**

**END OF PART A OF SECTION II**

**CALCULUS BC**  
**SECTION II, Part B**  
**Time—45 minutes**  
**Number of problems—3**

No calculator is allowed for these problems.



4. Let  $f$  be a function defined on the closed interval  $-5 \leq x \leq 5$  with  $f(1) = 3$ . The graph of  $f'$ , the derivative of  $f$ , consists of two semicircles and two line segments, as shown above.
- For  $-5 < x < 5$ , find all values  $x$  at which  $f$  has a relative maximum. Justify your answer.
  - For  $-5 < x < 5$ , find all values  $x$  at which the graph of  $f$  has a point of inflection. Justify your answer.
  - Find all intervals on which the graph of  $f$  is concave up and also has positive slope. Explain your reasoning.
  - Find the absolute minimum value of  $f(x)$  over the closed interval  $-5 \leq x \leq 5$ . Explain your reasoning.

**WRITE ALL WORK IN THE EXAM BOOKLET.**

5. Consider the differential equation  $\frac{dy}{dx} = 3x + 2y + 1$ .
- (a) Find  $\frac{d^2y}{dx^2}$  in terms of  $x$  and  $y$ .
- (b) Find the values of the constants  $m$ ,  $b$ , and  $r$  for which  $y = mx + b + e^{rx}$  is a solution to the differential equation.
- (c) Let  $y = f(x)$  be a particular solution to the differential equation with the initial condition  $f(0) = -2$ . Use Euler's method, starting at  $x = 0$  with a step size of  $\frac{1}{2}$ , to approximate  $f(1)$ . Show the work that leads to your answer.
- (d) Let  $y = g(x)$  be another solution to the differential equation with the initial condition  $g(0) = k$ , where  $k$  is a constant. Euler's method, starting at  $x = 0$  with a step size of 1, gives the approximation  $g(1) \approx 0$ . Find the value of  $k$ .
- 

6. Let  $f$  be the function given by  $f(x) = 6e^{-x/3}$  for all  $x$ .
- (a) Find the first four nonzero terms and the general term for the Taylor series for  $f$  about  $x = 0$ .
- (b) Let  $g$  be the function given by  $g(x) = \int_0^x f(t) dt$ . Find the first four nonzero terms and the general term for the Taylor series for  $g$  about  $x = 0$ .
- (c) The function  $h$  satisfies  $h(x) = kf'(ax)$  for all  $x$ , where  $a$  and  $k$  are constants. The Taylor series for  $h$  about  $x = 0$  is given by

$$h(x) = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + \cdots$$

Find the values of  $a$  and  $k$ .

---

**WRITE ALL WORK IN THE EXAM BOOKLET.**

**END OF EXAM**