



AP[®] Calculus BC 2009 Scoring Guidelines Form B

The College Board

The College Board is a not-for-profit membership association whose mission is to connect students to college success and opportunity. Founded in 1900, the association is composed of more than 5,600 schools, colleges, universities and other educational organizations. Each year, the College Board serves seven million students and their parents, 23,000 high schools and 3,800 colleges through major programs and services in college readiness, college admissions, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT[®], the PSAT/NMSQT[®] and the Advanced Placement Program[®] (AP[®]). The College Board is committed to the principles of excellence and equity, and that commitment is embodied in all of its programs, services, activities and concerns.

© 2009 The College Board. College Board, Advanced Placement Program, AP, AP Central, SAT, and the acorn logo are registered trademarks of the College Board. PSAT/NMSQT is a registered trademark of the College Board and National Merit Scholarship Corporation.


Permission to use copyrighted College Board materials may be requested online at:
www.collegeboard.com/inquiry/cbpermit.html.

Visit the College Board on the Web: www.collegeboard.com.
AP Central[®] is the official online home for AP teachers: apcentral.collegeboard.com.

AP[®] CALCULUS BC
2009 SCORING GUIDELINES (Form B)

Question 1

A baker is creating a birthday cake. The base of the cake is the region R in the first quadrant under the graph of $y = f(x)$ for $0 \leq x \leq 30$, where $f(x) = 20\sin\left(\frac{\pi x}{30}\right)$. Both x and y are measured in centimeters. The region R is shown in the figure above. The derivative of f is $f'(x) = \frac{2\pi}{3}\cos\left(\frac{\pi x}{30}\right)$.



- (a) The region R is cut out of a 30-centimeter-by-20-centimeter rectangular sheet of cardboard, and the remaining cardboard is discarded. Find the area of the discarded cardboard.
- (b) The cake is a solid with base R . Cross sections of the cake perpendicular to the x -axis are semicircles. If the baker uses 0.05 gram of unsweetened chocolate for each cubic centimeter of cake, how many grams of unsweetened chocolate will be in the cake?
- (c) Find the perimeter of the base of the cake.

(a) Area = $30 \cdot 20 - \int_0^{30} f(x) \, dx = 218.028 \text{ cm}^2$

3 : $\begin{cases} 2 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) Volume = $\int_0^{30} \frac{\pi}{2} \left(\frac{f(x)}{2}\right)^2 \, dx = 2356.194 \text{ cm}^3$

Therefore, the baker needs $2356.194 \times 0.05 = 117.809$ or 117.810 grams of chocolate.

3 : $\begin{cases} 2 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(c) Perimeter = $30 + \int_0^{30} \sqrt{1 + (f'(x))^2} \, dx = 81.803$ or 81.804 cm

3 : $\begin{cases} 2 : \text{integral} \\ 1 : \text{answer} \end{cases}$

AP[®] CALCULUS BC
2009 SCORING GUIDELINES (Form B)

Question 2

A storm washed away sand from a beach, causing the edge of the water to get closer to a nearby road. The rate at which the distance between the road and the edge of the water was changing during the storm is modeled by $f(t) = \sqrt{t} + \cos t - 3$ meters per hour, t hours after the storm began. The edge of the water was 35 meters from the road when the storm began, and the storm lasted 5 hours. The derivative of $f(t)$ is $f'(t) = \frac{1}{2\sqrt{t}} - \sin t$.

- What was the distance between the road and the edge of the water at the end of the storm?
- Using correct units, interpret the value $f'(4) = 1.007$ in terms of the distance between the road and the edge of the water.
- At what time during the 5 hours of the storm was the distance between the road and the edge of the water decreasing most rapidly? Justify your answer.
- After the storm, a machine pumped sand back onto the beach so that the distance between the road and the edge of the water was growing at a rate of $g(p)$ meters per day, where p is the number of days since pumping began. Write an equation involving an integral expression whose solution would give the number of days that sand must be pumped to restore the original distance between the road and the edge of the water.

(a) $35 + \int_0^5 f(t) dt = 26.494$ or 26.495 meters

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

(b) Four hours after the storm began, the rate of change of the distance between the road and the edge of the water is increasing at a rate of 1.007 meters/hours².

$$2 : \begin{cases} 1 : \text{interpretation of } f'(4) \\ 1 : \text{units} \end{cases}$$

(c) $f'(t) = 0$ when $t = 0.66187$ and $t = 2.84038$
The minimum of f for $0 \leq t \leq 5$ may occur at 0, 0.66187, 2.84038, or 5.

$$3 : \begin{cases} 1 : \text{considers } f'(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$$

$$f(0) = -2$$

$$f(0.66187) = -1.39760$$

$$f(2.84038) = -2.26963$$

$$f(5) = -0.48027$$


The distance between the road and the edge of the water was decreasing most rapidly at time $t = 2.840$ hours after the storm began.

(d) $-\int_0^5 f(t) dt = \int_0^x g(p) dp$

$$2 : \begin{cases} 1 : \text{integral of } g \\ 1 : \text{answer} \end{cases}$$

AP[®] CALCULUS BC
2009 SCORING GUIDELINES (Form B)

Question 3



Graph of f

A continuous function f is defined on the closed interval $-4 \leq x \leq 6$. The graph of f consists of a line segment and a curve that is tangent to the x -axis at $x = 3$, as shown in the figure above. On the interval $0 < x < 6$, the function f is twice differentiable, with $f''(x) > 0$.

- (a) Is f differentiable at $x = 0$? Use the definition of the derivative with one-sided limits to justify your answer.
- (b) For how many values of a , $-4 \leq a < 6$, is the average rate of change of f on the interval $[a, 6]$ equal to 0? Give a reason for your answer.
- (c) Is there a value of a , $-4 \leq a < 6$, for which the Mean Value Theorem, applied to the interval $[a, 6]$, guarantees a value c , $a < c < 6$, at which $f'(c) = \frac{1}{3}$? Justify your answer.
- (d) The function g is defined by $g(x) = \int_0^x f(t) dt$ for $-4 \leq x \leq 6$. On what intervals contained in $[-4, 6]$ is the graph of g concave up? Explain your reasoning.

(a)
$$\lim_{h \rightarrow 0^-} \frac{f(h) - f(0)}{h} = \frac{2}{3}$$

$$\lim_{h \rightarrow 0^+} \frac{f(h) - f(0)}{h} < 0$$

Since the one-sided limits do not agree, f is not differentiable at $x = 0$.

(b)
$$\frac{f(6) - f(a)}{6 - a} = 0 \text{ when } f(a) = f(6).$$
 There are two values of a for which this is true.

(c) Yes, $a = 3$. The function f is differentiable on the interval $3 < x < 6$ and continuous on $3 \leq x \leq 6$.

Also,
$$\frac{f(6) - f(3)}{6 - 3} = \frac{1 - 0}{6 - 3} = \frac{1}{3}.$$

By the Mean Value Theorem, there is a value c ,

$$3 < c < 6, \text{ such that } f'(c) = \frac{1}{3}.$$

(d)
$$g'(x) = f(x), \quad g''(x) = f'(x)$$

$$g''(x) > 0 \text{ when } f'(x) > 0$$

This is true for $-4 < x < 0$ and $3 < x < 6$.

2 : $\begin{cases} 1 : \text{sets up difference quotient at } x = 0 \\ 1 : \text{answer with justification} \end{cases}$

2 : $\begin{cases} 1 : \text{expression for average rate of change} \\ 1 : \text{answer with reason} \end{cases}$


2 : $\begin{cases} 1 : \text{answers "yes" and identifies } a = 3 \\ 1 : \text{justification} \end{cases}$

3 : $\begin{cases} 1 : g'(x) = f(x) \\ 1 : \text{considers } g''(x) > 0 \\ 1 : \text{answer} \end{cases}$

AP[®] CALCULUS BC
2009 SCORING GUIDELINES (Form B)

Question 4

The graph of the polar curve $r = 1 - 2\cos \theta$ for $0 \leq \theta \leq \pi$ is shown above. Let S be the shaded region in the third quadrant bounded by the curve and the x -axis.



- (a) Write an integral expression for the area of S .
- (b) Write expressions for $\frac{dx}{d\theta}$ and $\frac{dy}{d\theta}$ in terms of θ .
- (c) Write an equation in terms of x and y for the line tangent to the graph of the polar curve at the point where $\theta = \frac{\pi}{2}$.

Show the computations that lead to your answer.

(a) $r(0) = -1$; $r(\theta) = 0$ when $\theta = \frac{\pi}{3}$.

$$\text{Area of } S = \frac{1}{2} \int_0^{\pi/3} (1 - 2\cos \theta)^2 d\theta$$

2 : $\left\{ \begin{array}{l} 1 : \text{limits and constant} \\ 1 : \text{integrand} \end{array} \right.$

(b) $x = r\cos \theta$ and $y = r\sin \theta$

$$\frac{dr}{d\theta} = 2\sin \theta$$

$$\frac{dx}{d\theta} = \frac{dr}{d\theta} \cos \theta - r \sin \theta = 4\sin \theta \cos \theta - \sin \theta$$

$$\frac{dy}{d\theta} = \frac{dr}{d\theta} \sin \theta + r \cos \theta = 2\sin^2 \theta + (1 - 2\cos \theta) \cos \theta$$

4 : $\left\{ \begin{array}{l} 1 : \text{uses } x = r\cos \theta \text{ and } y = r\sin \theta \\ 1 : \frac{dr}{d\theta} \\ 2 : \text{answer} \end{array} \right.$

(c) When $\theta = \frac{\pi}{2}$, we have $x = 0$, $y = 1$.

$$\left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{2}} = \left. \frac{dy/d\theta}{dx/d\theta} \right|_{\theta=\frac{\pi}{2}} = -2$$


The tangent line is given by $y = 1 - 2x$.

3 : $\left\{ \begin{array}{l} 1 : \text{values for } x \text{ and } y \\ 1 : \text{expression for } \frac{dy}{dx} \\ 1 : \text{tangent line equation} \end{array} \right.$

AP[®] CALCULUS BC
2009 SCORING GUIDELINES (Form B)

Question 5

Let f be a twice-differentiable function defined on the interval $-1.2 < x < 3.2$ with $f(1) = 2$. The graph of f' , the derivative of f , is shown above. The graph of f' crosses the x -axis at $x = -1$ and $x = 3$ and has a horizontal tangent at $x = 2$. Let g be the function given by $g(x) = e^{f(x)}$.



- (a) Write an equation for the line tangent to the graph of g at $x = 1$.
- (b) For $-1.2 < x < 3.2$, find all values of x at which g has a local maximum. Justify your answer.
- (c) The second derivative of g is $g''(x) = e^{f(x)}[(f'(x))^2 + f''(x)]$. Is $g''(-1)$ positive, negative, or zero? Justify your answer.
- (d) Find the average rate of change of g' , the derivative of g , over the interval $[1, 3]$.

(a) $g(1) = e^{f(1)} = e^2$
 $g'(x) = e^{f(x)} f'(x)$, $g'(1) = e^{f(1)} f'(1) = -4e^2$
 The tangent line is given by $y = e^2 - 4e^2(x - 1)$.

$$3 : \begin{cases} 1 : g'(x) \\ 1 : g(1) \text{ and } g'(1) \\ 1 : \text{tangent line equation} \end{cases}$$

(b) $g'(x) = e^{f(x)} f'(x)$
 $e^{f(x)} > 0$ for all x
 So, g' changes from positive to negative only when f' changes from positive to negative. This occurs at $x = -1$ only. Thus, g has a local maximum at $x = -1$.

$$2 : \begin{cases} 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$$

(c) $g''(-1) = e^{f(-1)}[(f'(-1))^2 + f''(-1)]$
 $e^{f(-1)} > 0$ and $f'(-1) = 0$
 Since f' is decreasing on a neighborhood of -1 , $f''(-1) < 0$. Therefore, $g''(-1) < 0$.

$$2 : \begin{cases} 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$$

(d) $\frac{g'(3) - g'(1)}{3 - 1} = \frac{e^{f(3)} f'(3) - e^{f(1)} f'(1)}{2} = 2e^2$

$$2 : \begin{cases} 1 : \text{difference quotient} \\ 1 : \text{answer} \end{cases}$$

AP[®] CALCULUS BC
2009 SCORING GUIDELINES (Form B)

Question 6

The function f is defined by the power series

$$f(x) = 1 + (x + 1) + (x + 1)^2 + \cdots + (x + 1)^n + \cdots = \sum_{n=0}^{\infty} (x + 1)^n$$

for all real numbers x for which the series converges.

- (a) Find the interval of convergence of the power series for f . Justify your answer.
- (b) The power series above is the Taylor series for f about $x = -1$. Find the sum of the series for f .
- (c) Let g be the function defined by $g(x) = \int_{-1}^x f(t) dt$. Find the value of $g\left(-\frac{1}{2}\right)$, if it exists, or explain why $g\left(-\frac{1}{2}\right)$ cannot be determined.
- (d) Let h be the function defined by $h(x) = f(x^2 - 1)$. Find the first three nonzero terms and the general term of the Taylor series for h about $x = 0$, and find the value of $h\left(\frac{1}{2}\right)$.

- (a) The power series is geometric with ratio $(x + 1)$.
The series converges if and only if $|x + 1| < 1$.
Therefore, the interval of convergence is $-2 < x < 0$.

OR

$$\lim_{n \rightarrow \infty} \left| \frac{(x + 1)^{n+1}}{(x + 1)^n} \right| = |x + 1| < 1 \text{ when } -2 < x < 0$$

At $x = -2$, the series is $\sum_{n=0}^{\infty} (-1)^n$, which diverges since the

terms do not converge to 0. At $x = 0$, the series is $\sum_{n=0}^{\infty} 1$,

which similarly diverges. Therefore, the interval of convergence is $-2 < x < 0$.

- (b) Since the series is geometric,

$$f(x) = \sum_{n=0}^{\infty} (x + 1)^n = \frac{1}{1 - (x + 1)} = -\frac{1}{x} \text{ for } -2 < x < 0.$$

- (c) $g\left(-\frac{1}{2}\right) = \int_{-1}^{-\frac{1}{2}} -\frac{1}{x} dx = -\ln|x| \Big|_{x=-1}^{x=-\frac{1}{2}} = \ln 2$

- (d) $h(x) = f(x^2 - 1) = 1 + x^2 + x^4 + \cdots + x^{2n} + \cdots$

$$h\left(\frac{1}{2}\right) = f\left(-\frac{3}{4}\right) = \frac{4}{3}$$

$$3 : \begin{cases} 1 : \text{identifies as geometric} \\ 1 : |x + 1| < 1 \\ 1 : \text{interval of convergence} \end{cases}$$

OR

$$3 : \begin{cases} 1 : \text{sets up limit of ratio} \\ 1 : \text{radius of convergence} \\ 1 : \text{interval of convergence} \end{cases}$$

1 : answer

$$2 : \begin{cases} 1 : \text{antiderivative} \\ 1 : \text{value} \end{cases}$$

$$3 : \begin{cases} 1 : \text{first three terms} \\ 1 : \text{general term} \\ 1 : \text{value of } h\left(\frac{1}{2}\right) \end{cases}$$