

AP[®] Calculus AB Exam

SECTION II: Free Response

2017

DO NOT OPEN THIS BOOKLET OR BREAK THE SEALS ON PART B UNTIL YOU ARE TOLD TO DO SO.

At a Glance

Total Time

1 hour and 30 minutes

Number of Questions

6

Percent of Total Score

50%

Writing Instrument

Either pencil or pen with black or dark blue ink

Weight

The questions are weighted equally, but the parts of a question are not necessarily given equal weight.

Part A

Number of Questions

2

Time

30 minutes

Electronic Device

Graphing calculator required

Percent of Section II Score

33.33%

Part B

Number of Questions

4

Time

1 hour

Electronic Device

None allowed

Percent of Section II Score

66.67%

IMPORTANT Identification Information

PLEASE PRINT WITH PEN:

1. First two letters of your last name
First letter of your first name
2. Date of birth

Month Day Year
3. Six-digit school code
4. Unless I check the box below, I grant the College Board the unlimited right to use, reproduce, and publish my free-response materials, both written and oral, for educational research and instructional purposes. My name and the name of my school will not be used in any way in connection with my free-response materials. I understand that I am free to mark "No" with no effect on my score or its reporting.
No, I do not grant the College Board these rights.

Instructions

The questions for Section II are printed in this booklet. Do not break the seals on Part B until you are told to do so. Write your solution to each part of each question in the space provided. Write clearly and legibly. Cross out any errors you make; erased or crossed-out work will not be scored.

Manage your time carefully. During Part A, work only on the questions in Part A. You are permitted to use your calculator to solve an equation, find the derivative of a function at a point, or calculate the value of a definite integral. However, you must clearly indicate the setup of your question, namely the equation, function, or integral you are using. If you use other built-in features or programs, you must show the mathematical steps necessary to produce your results. During Part B, you may continue to work on the questions in Part A without the use of a calculator.

As you begin each part, you may wish to look over the questions before starting to work on them. It is not expected that everyone will be able to complete all parts of all questions.

- Show all of your work, even though a question may not explicitly remind you to do so. Clearly label any functions, graphs, tables, or other objects that you use. Justifications require that you give mathematical reasons, and that you verify the needed conditions under which relevant theorems, properties, definitions, or tests are applied. Your work will be scored on the correctness and completeness of your methods as well as your answers. Answers without supporting work will usually not receive credit.
- Your work must be expressed in standard mathematical notation rather than calculator syntax. For example, $\int_1^5 x^2 dx$ may not be written as `fnInt(X2, X, 1, 5)`.
- Unless otherwise specified, answers (numeric or algebraic) need not be simplified. If you use decimal approximations in calculations, your work will be scored on accuracy. Unless otherwise specified, your final answers should be accurate to three places after the decimal point.
- Unless otherwise specified, the domain of a function f is assumed to be the set of all real numbers x for which $f(x)$ is a real number.

Form I

Form Code 4NBP4-S

66

CALCULUS AB
SECTION II, Part A
Time—30 minutes
Number of questions—2

A GRAPHING CALCULATOR IS REQUIRED FOR THESE QUESTIONS.

1**1****1****1****1****1****1****1****1****1**

1. A particle moves along the x -axis so that its velocity at time t is given by $v(t) = \frac{t^6 - 13t^4 + 12}{10t^3 + 3}$. At time

$t = 0$, the initial position of the particle is $x = 7$.

(a) Find the acceleration of the particle at time $t = 5.1$.

(b) Find all values of t in the interval $0 \leq t \leq 2$ for which the speed of the particle is 1.

Do not write beyond this border.

Do not write beyond this border.


1**1****1****1****1****1****1****1****1****1**

- (c) Find the position of the particle at time $t = 4$. Is the particle moving toward the origin or away from the origin at time $t = 4$? Justify your answer.

-
- (d) During the time interval $0 < t \leq 4$, does the particle return to its initial position? Give a reason for your answer.

Do not write beyond this border.

Do not write beyond this border.



2. During the time interval $0 \leq t \leq 4.5$ hours, water flows into tank A at a rate of $a(t) = (2t - 5) + 5e^{2\sin t}$ liters per hour. During the same time interval, water flows into tank B at a rate of $b(t)$ liters per hour. Both tanks are empty at time $t = 0$. The graphs of $y = a(t)$ and $y = b(t)$, shown in the figure above, intersect at $t = k$ and $t = 2.416$.

(a) How much water will be in tank A at time $t = 4.5$?

-
- (b) During the time interval $0 \leq t \leq k$ hours, water flows into tank B at a constant rate of 20.5 liters per hour. What is the difference between the amount of water in tank A and the amount of water in tank B at time $t = k$?

Do not write beyond this border.

Do not write beyond this border.

- (c) The area of the region bounded by the graphs of $y = a(t)$ and $y = b(t)$ for $k \leq t \leq 2.416$ is 14.470. How much water is in tank B at time $t = 2.416$?

-
- (d) During the time interval $2.7 \leq t \leq 4.5$ hours, the rate at which water flows into tank B is modeled by

$$w(t) = 21 - \frac{30t}{(t-8)^2} \text{ liters per hour. Is the difference } w(t) - a(t) \text{ increasing or decreasing at time}$$

$t = 3.5$? Show the work that leads to your answer.

Do not write beyond this border.

Do not write beyond this border.

END OF PART A

**IF YOU FINISH BEFORE TIME IS CALLED,
YOU MAY CHECK YOUR WORK ON PART A ONLY.**


DO NOT GO ON TO PART B UNTIL YOU ARE TOLD TO DO SO.

CALCULUS AB
SECTION II, Part B
Time—1 hour
Number of questions—4

NO CALCULATOR IS ALLOWED FOR THESE QUESTIONS.

DO NOT BREAK THE SEALS UNTIL YOU ARE TOLD TO DO SO.

NO CALCULATOR ALLOWED

Graph of f

3. The graph of the function f , consisting of three line segments and a quarter of a circle, is shown above. Let g

be the function defined by $g(x) = \int_1^x f(t) dt$.

- (a) Find the average rate of change of g from $x = -5$ to $x = 5$.

-
- (b) Find the instantaneous rate of change of g with respect to x at $x = 3$, or state that it does not exist.

Do not write beyond this border.

Do not write beyond this border.

3

3

3

3

3

3

3

3

3

3

NO CALCULATOR ALLOWED

(c) On what open intervals, if any, is the graph of g concave up? Justify your answer.

(d) Find all x -values in the interval $-5 < x < 5$ at which g has a critical point. Classify each critical point as the location of a local minimum, a local maximum, or neither. Justify your answers.

Do not write beyond this border.

Do not write beyond this border.

NO CALCULATOR ALLOWED

x	0	1	2	3	4	5	6
$f'(x)$	4	3.5	2	0.8	1.7	5.8	7

4. The function f satisfies $f(0) = 20$. The first derivative of f satisfies the inequality $0 \leq f'(x) \leq 7$ for all x in the closed interval $[0, 6]$. Selected values of f' are shown in the table above. The function f has a continuous second derivative for all real numbers.

(a) Use a midpoint Riemann sum with three subintervals of equal length indicated by the data in the table to approximate the value of $f(6)$.

(b) Determine whether the actual value of $f(6)$ could be 70. Explain your reasoning.

Do not write beyond this border.

Do not write beyond this border.

4

4

4

4

4

4

4

4

4

4

NO CALCULATOR ALLOWED


(c) Evaluate $\int_2^4 f''(x) dx$.

(d) Find $\lim_{x \rightarrow 0} \frac{f(x) - 20e^x}{0.5f(x) - 10}$.

Do not write beyond this border.

Do not write beyond this border.

NO CALCULATOR ALLOWED



5. Let R be the region in the first quadrant enclosed by the graph of $f(x) = \sqrt{\cos x}$, the graph of $g(x) = e^x$, and the vertical line $x = \frac{\pi}{2}$, as shown in the figure above.

(a) Write, but do not evaluate, an integral expression that gives the area of R .

Do not write beyond this border.

Do not write beyond this border.

5

5

5

5

5

5

5

5

5

5

NO CALCULATOR ALLOWED

(b) Find the volume of the solid generated when R is revolved about the x -axis.

(c) Region R is the base of a solid whose cross sections perpendicular to the x -axis are semicircles with diameters on the xy -plane. Write, but do not evaluate, an integral expression that gives the volume of this solid.

Do not write beyond this border.

Do not write beyond this border.

6

6

6

6

6

6

6

6

6

6

NO CALCULATOR ALLOWED

6. For $0 \leq t \leq 6$ seconds, a screen saver on a computer screen shows two circles that start as dots and expand outward.
- (a) At the instant that the first circle has a radius of 9 centimeters, the radius is increasing at a rate of $\frac{3}{2}$ centimeters per second. Find the rate at which the area of the circle is changing at that instant. Indicate units of measure.

-
- (b) The radius of the first circle is modeled by $w(t) = 12 - 12e^{-0.5t}$ for $0 \leq t \leq 6$, where $w(t)$ is measured in centimeters and t is measured in seconds. At what time t is the radius of the circle increasing at a rate of 3 centimeters per second?

Do not write beyond this border.

Do not write beyond this border.

6**6****6****6****6****6****6****6****6****6****NO CALCULATOR ALLOWED**

- (c) A model for the radius of the second circle is given by the function f for $0 \leq t \leq 6$, where $f(t)$ is measured in centimeters and t is measured in seconds. The rate of change of the radius of the second circle is given by $f'(t) = t^2 - 4t + 4$. Based on this model, by how many centimeters does the radius of the second circle increase from time $t = 0$ to $t = 3$?

Do not write beyond this border.

Do not write beyond this border.